

HBT Interferometry with



Frank Laue, *The Ohio State University*
for the STAR Collaboration


- Au+Au Collisions at STAR
- **A Technical Issue**
 - Track Merging
- **3D -HBT**
 - Coulomb Correction
- **Bread and Butter ... - Pion HBT**
 - Centrality
 - Transverse Momentum (p_T, k_T)
 - The HBT Excitation Function
- **... and beyond**
 - E-by-E HBT
 - Pion Phase Space Density
 - K_S^0
- Summary
- Future Plans



Au+Au Collisions as seen from STAR

- More than 2M events were taken in Y2K at $\sqrt{s_{NN}} = 130\text{GeV}$

Front view of the TPC




Particle identification via specific ionisation (dE/dx):

Optimum performance for HBT:

$$-0.5 < y < +0.5$$

$$0.125 < p_T / (\text{GeV}/c) < 0.450$$

π : electron contamination < 10%




Two Particle Interferometry

Two particle correlations are sensitive space-time geometry

$$C(p_1, p_2) = \frac{P(p_1, p_2)}{P(p_1)P(p_2)} = \frac{N_{real}}{N_{mixed}} = 1 + |\tilde{\rho}(q)|^2$$

$\pi\pi : |y| < 0.5$

$0.125 < p_T(\text{GeV}/c) < 0.225$



The simplest parameterization:

$$C(Q_{inv}) = 1 + \lambda \cdot e^{-Q_{inv}^2 R_{inv}^2}$$


$$Q_{inv} = \sqrt{(p_1 - p_2)^2 - (E_1 - E_2)^2}$$

$$R_{inv} = (6.3 \pm 0.5) \text{ fm}$$


$$\lambda = 0.5 \pm 0.1$$



A Technical Issue: Track Merging



Close tracks can be reconstructed as one
→ artificial suppression of low- q pairs




- Cutting on the entrance separation of the TPC removes the merging effect, but de-emphasizes low- q pairs.
- For non-Gaussian sources this leads to a residual effect on the HBT parameters requiring correction $\sim 10\%$ in λ and $\sim 4\%$ in the radii.

3Dimensional - HBT

Pratt-Bertsch Parameterization (measured in the LCMS frame; $(\mathbf{p}_1 + \mathbf{p}_2)_z = 0$)

$$C(q_{Out}, q_{Side}, q_{Long}) = 1 + \lambda e^{-(q_{Out}^2 R_{Out}^2 + q_{Side}^2 R_{Side}^2 + q_{Long}^2 R_{Long}^2)}$$




Information:

- geometrical source size

- lifetime $\tau = \sqrt{R_{Out}^2 - R_{Side}^2} / \beta_T$

(for transparent sources)




Soff et al. nucl-th/0012085



$$K_T = (\mathbf{p}_1 + \mathbf{p}_2)_T / 2$$

3 Dimensional - $\pi^- \pi^-$ HBT

1D Projections of the 3D Pratt-Bertsch Parameterization



12% most central events

Pion tracks:

$0.125 < p_T / (\text{GeV/c}) < 0.225$;
 $|y| < 0.5$

- Coulomb corrected with a 5 fm source
- No Coulomb correction

$$\lambda = 0.50 \pm 0.01 \pm 0.03$$

$$R_{Out} = (5.86 \pm 0.11 \pm 0.23) \text{ fm}$$

$$R_{Side} = (5.47 \pm 0.09 \pm 0.16) \text{ fm}$$

$$R_{Long} = (7.07 \pm 0.12 \pm 0.21) \text{ fm}$$

sys. error result from the pair cuts (merging)
 and Coulomb correction



Coulomb Correction


Standard Coulomb

5-dimensional Monte-Carlo integration of Coulomb wave functions over a spherical source {S.Pratt et al., Phys.Rev. C42(1990)2646}


Damped Coulomb ($\lambda < 1$)

- particle misidentification
- long lived resonances

Size dependence



Damping dependence



Centrality Dependence

$0.125 \text{ GeV}/c < p_T < 0.225 \text{ GeV}/c$


• π^- • π^+

STAR
Preliminary

- $\pi^+\pi^+$, $\pi^-\pi^-$, parameters similar
- λ 's don't change with mult.
- radii increase with multiplicity
- roughly similar to AGS/SPS

- Comparison with AGS/SPS
- parameters roughly same
 - similar increase in R_{Out} , R_{Side} (geometric effect)

- R_{Long} - increase not seen at AGS/SPS



m_T Dependence - sensitivity to flow

12 % most central events


• π^- • π^+

STAR
Preliminary

- $\pi^+\pi^+$, $\pi^-\pi^-$ HBT parameters similar
- With increasing m_T
 - λ increases \rightarrow fewer resonances
 - radii decrease \rightarrow position-momentum correlations


- Comparison with AGS/SPS
- parameters roughly similar to AGS/SPS
 - With increasing m_T
 - similar increase in λ
 - similar decrease of radii

- stronger effect in R_{Out} than at AGS/SPS




The R_{Out}/R_{Side} Ratio

emission duration for transparent sources: $\tau = \sqrt{R_{Out}^2 - R_{Side}^2} / \beta_T$



$$K_T = (p_T^1 + p_T^2) / 2$$

Tomášik, Heinz nucl-th/9805016



x-p correlations (flow / opaqueness) might change the homogeneity length

Note: extremely strong flow at RHIC (close to hydro)




The HBT Excitation function

Compilation of world BP
3D $\pi\pi$ -HBT parameters as
a function of \sqrt{s}



- ~10% Central AuAu(PbPb) events
- $y \sim 0$
- $k_T \approx 0.17$ GeV/c


no significant rise in spatio-temporal
size of the π emitting source



The π Phase Space Density (Poster by J.Cramer)

pion occupation of cell in
coordinate \otimes momentum space:


$$\langle f \rangle(m_T) = \frac{(\hbar c)^3}{2 E_x m_T} \frac{d^2 N}{dy dm_T} \frac{(\lambda \pi)^{1/2}}{R_O R_S R_L}$$



- "Universal" phase space density observed at SPS appears to hold at RHIC as well
- Consistent with thermal distribution ($T=94\text{MeV}$) and strong collective flow ($\beta = 0.58$)
- Fundamental phase space saturation may relate increases in geometry, temperature, multiplicity




m_T systematics ($\pi\pi, \bar{d}/\bar{p}$)



\bar{d}/\bar{p} - Talk by D.Hardtke





Event by Event - HBT (Poster by D.Flierl)




- Numerator from one event only
- Denominator from 10 mixed events
- Numerators and Denominators from $\pi^+ \pi^+$ and $\pi^- \pi^-$ HBT added
- for 50% of all events the fit fails ($0 < \lambda < 1$)
 - these events are excluded from further analysis

Event characterization EbyE ? almost




$K_S^0 K_S^0$ - Correlations (Poster by T.Humanic)

1.3 K^0 /event



- no coulomb repulsion problems
 - less 2 track resolution problems
 - few distortions from resonance decays
- ideal for HBT



K_S^0 Correlation will become statistically meaningful once we have 10M events



Summary

- first $\pi\pi$ interferometry at RHIC
 - source sizes roughly same as at AGS/SPS ($< 10\text{fm}$)
 - radii increase with centrality (expected for $R_{\text{Out}}, R_{\text{Side}}$)
 - radii decrease with increasing k_T (flow)
 - $R_{\text{Out}}/R_{\text{Side}} \sim 1$ (explosive source)
 - The “universal” phase space density observed at SPS holds at RHIC
- promising novel analyses
 - event characterisation E-by-E: almost
 - K^0 interferometry

Future Plans

- improved Coulomb correction
- different collision systems
- other particle combinations
- **energy scan**



π^+	π^-	K^+	K^-	K^0	p	\bar{p}	Λ	$\bar{\Lambda}$	
█	█	█	█	█	█ Δ^+	█	█	█	π^+
	█	█	█	█	█ Δ^0	█	█	█	π^-
		█	█	█	█	█	█	█	K^+
			█	█	█	█	█	█	K^-
				█	█	█	█	█	K^0
					█	█	█	█	p
						█	█	█	\bar{p}
							█	█	Λ
								█	$\bar{\Lambda}$

The next steps:
Exploring the HBT matrix

- $K^0K^0, \Lambda p, \Lambda\Lambda$ T. Humanic
- event-by-event HBT radii D. Flierl
- HBT vs. reaction plane R. Wells
- $\bar{p}\bar{p}, p\bar{p}$ M. Lopez-Noriega
- 6-D phase space density J. Cramer
- $K^+\pi^-$ (emission-order) A. Kiesel
- 3- π correlations R. Willson

“traditional”
HBT axis